





Condensation in 2D: from non-universal effects to bubble traps

Andrea Tononi, Alberto Cappellaro, Luca Salasnich

Dipartimento di Fisica e Astronomia "Galileo Galilei" and CNISM,
Università di Padova, Via Marzolo 8, I-35131 Padova, Italy




Graphical abstract




Nonuniversal effects

Bubble Trap



from N. Lundblad et al., arXiv:1906.05885v1



Functional integration

We consider a uniform D -dimensional Bose gas of identical cold atoms with mass m , described by the complex field $\psi(\vec{r}, \tau)$.

The partition function

$$\mathcal{Z} = \int D(\bar{\psi}, \psi) e^{-\frac{S[\bar{\psi}, \psi]}{\hbar}},$$

$$S[\bar{\psi}, \psi] = \int_0^{\beta\hbar} d\tau \int d^D r \mathcal{L},$$

$$\mathcal{L} = \bar{\psi}_{\vec{r}, \tau} \left(\hbar \partial_\tau - \frac{\hbar^2 \nabla^2}{2m} - \mu \right) \psi_{\vec{r}, \tau} + \frac{1}{2} \int d^D r' |\psi_{\vec{r}, \tau}|^2 V(\vec{r} - \vec{r}') |\psi_{\vec{r}', \tau}|^2$$

Performing functional integration at a Gaussian level ($\psi_{\vec{r}, \tau} = \psi_0 + \eta_{\vec{r}, \tau}$), one obtains $\Omega = -\beta^{-1} \ln \mathcal{Z}$ as

$$\Omega(\mu, \psi_0^2, T) = \Omega_0 + \Omega_g^{(0)} + \Omega_g^{(T)}$$

with the excitation spectrum

$$E_{\vec{k}} = \sqrt{(\varepsilon_{\vec{k}} - \mu + g_0 \psi_0^2 + \psi_0^2 \tilde{V}(\vec{k}))^2 - (\psi_0^2 \tilde{V}(\vec{k}))^2}$$

From nonuniversal effects to bubble traps

Nonuniversal effects Weakly-interacting bosons are usually described with the zero-range interaction $\tilde{V}(\mathbf{k}) = g_0$. With scattering theory, g_0 is linked to the s-wave scattering length a_s (universal parameter). We study $\tilde{V}(\mathbf{k}) = g_0 + g_2 k^2$, including the first nonzero correction in the momentum. Here g_0 and g_2 are linked to a_s and:

- r_{eff} in 3D (effective range)
- R_{ch} in 2D (characteristic range)

In specific regimes (in 3D: $a_s/r_{\text{eff}} \leq 1$, in 2D: $a_s/R_{\text{ch}} \leq 1$) the finite-range corrections to the thermodynamics are relevant.

Bubble traps Bubble traps are produced with the external potential

$$U(\vec{r}) = \sqrt{(u(\vec{r}) + \hbar \Delta)^2 + \hbar \omega^2},$$

with $u(\vec{r})$ an harmonic potential. In the thin-shell limit, we simply put $V = S^2$.

Bibliography

- [1] A. Tononi, A. Cappellaro, L. Salasnich, New J. of Phys., **20**, 125007 (2018).
- [2] A. Tononi, Condens. Matter **4**, 20 (2019).
- [3] A. Tononi, L. Salasnich, arXiv:1903.08453 (2019).

Nonuniversal equation of state in 2D

With the superfluid parametrization $\psi = \sqrt{\rho_0 + \delta \rho} e^{i\theta}$, we derive in Ref. [2] the zero-temperature equation of state for a 2D homogeneous Bose gas

$$P(\mu, T=0) = \frac{m\mu^2}{8\pi\hbar^2\lambda^{3/2}} \left[\ln\left(\frac{\epsilon_0}{\mu}\lambda\right) - \frac{1}{2} \right], \quad \lambda = 1 + \frac{4m\mu}{\hbar^2 g_0}$$

For $na_s^2 = 10^{-5}$ and $nR_{\text{ch}}^2 = 6 \times 10^{-2}$ we get a **20%** correction.

BEC on a sphere

In Ref. [3], we apply VPT to calculate the critical temperature T_{BEC} for interacting bosons confined on a 2D spherical shell

$$k_B T_{\text{BEC}} = \left(\frac{2\pi\hbar^2 n}{m} - \frac{gn}{2} \right) / \left[\frac{\hbar^2 \beta_{\text{BEC}}}{2mR^2} \left(1 + \sqrt{1 + \frac{2gmnR^2}{\hbar^2}} \right) - \ln \left(e^{\frac{\hbar^2 \beta_{\text{BEC}}}{mR^2} \sqrt{1 + \frac{2gmnR^2}{\hbar^2}}} - 1 \right) \right],$$

and the condensate fraction



$$\frac{n_0}{n} = 1 - \frac{mg}{4\pi\hbar^2} - \frac{1}{4\pi R^2 n} \left[1 + \sqrt{1 + \frac{2gmnR^2}{\hbar^2}} \right] + \frac{mk_B T}{2\pi\hbar^2 n} \times$$

$$\ln \left(e^{\frac{\hbar^2}{mR^2 k_B T} \sqrt{1 + \frac{2gmnR^2}{\hbar^2}}} - 1 \right).$$


In the limit $R \rightarrow \infty$ one has $T_{\text{BEC}} \rightarrow 0$, and n_0/n gives Schick result (PRA,3,1067).

We calculate n_s in analogy to the Landau formula, and obtain T_{BKT} applying the Kosterlitz-Nelson criterion, unchanged with respect to the flat plane (PRD,43,1314).

The static and dynamic properties of bosons in bubble-traps will be studied in the ISS (microgravity conditions).



Condensation and superfluidity in D dimensions




For the finite-range effective interaction $\tilde{V}(\mathbf{k}) = g_0 + g_2 k^2$ in Ref. [1] we calculate

- the number density n as a function of condensate density n_0 and T as

$$n(n_0, T) = -\frac{1}{L^D} \frac{\partial \Omega(\mu_e, \psi_0, T)}{\partial \mu_e} = n_0 + f_g^{(0)} + f_g^{(T)}$$


- the superfluid density $n_s = n(n_0, T) - n_n(n_0, T)$, with n_n the normal density given by a self-consistent derivation of Landau formula

Finite-range corrections to n_0/n



2D condensate fraction n_0/n at $T=0$ and $R_{\text{ch}} = 2a_s$, in terms of the gas parameter na_s^2 .

Finite-range corrections to n_s/n



3D superfluid fraction n_s/n for $r_{\text{eff}} = a_s$, as a function of the temperature $T^* = k_B T/E_r$, where $E_r = \hbar^2 n^{2/3}/m$.

Contact informations

Andrea Tononi
andrea.tononi@phd.unipd.it
www.andreatononi.com