

Binding of heavy fermions by a single light atom in 1D

Andrea Tononi

Laboratoire Physique Théorique et Modèles Statistiques,
CNRS, Université Paris-Saclay

ECT* NPES Workshop,
Trento


04-08/07/2022

based on

[A. Tononi, J. Givois, and D. S. Petrov, arXiv:2205.01018,
accepted in PRA as a Letter]

The system

$$\hat{H} = \int \left(-\frac{\hat{\Psi}_x^\dagger \partial_x^2 \hat{\Psi}_x}{2M} - \frac{\hat{\phi}_x^\dagger \partial_x^2 \hat{\phi}_x}{2m} + g \hat{\Psi}_x^\dagger \hat{\phi}_x^\dagger \hat{\Psi}_x \hat{\phi}_x \right) dx, \quad g < 0$$




N heavy fermions of mass M
1 light atom of mass m




Noninteracting heavy

heavy-light attraction

How many heavy fermions can be bound by a single light atom in 1D?




How many heavy fermions can be bound by a single light atom in 1D?



Competition between:

- ▶ Kinetic energy of heavy atoms $\sim 1/M$
- ▶ (Effective) attractive heavy-heavy potential, mediated by the exchange of the light atom $\sim 1/m$


The $(N+1)$ -body problem



A **well posed** problem (clear question), with few **simple** parameters

- ▶ spatial dimension $D = 1$,
- ▶ mass ratio M/m ,
- ▶ scattering length a
- ▶ number of heavy atoms N



The $(N+1)$ -body problem



A **well posed** problem (clear question), with few **simple** parameters

- ▶ spatial dimension $D = 1$,
- ▶ mass ratio M/m ,
- ▶ scattering length a
- ▶ number of heavy atoms N

relevant for **experiments with mass and density-imbalanced fermionic mixtures**




[Ravensbergen et al, PRA **98**, 063624 (2018)]

Outline

- ▷ Introduction and motivation
- ▷ Born-Oppenheimer theory of the 3D trimer
- ▷ Bound states of $N + 1$ fermions in 1D
- ▷ Derivation of the results
 - ▷ Exact results for $N \leq 5$
 - ▷ Mean field: Thomas-Fermi approximation
 - ▷ Mean field: Hartree-Fock
- ▷ Conclusions and perspectives

2+1 (trimer) in the 3D Born-Oppenheimer picture




Light atom in the field of fixed
heavy fermions

(distance $R = |\vec{R}_2 - \vec{R}_1|$)

$$-\frac{\hbar^2 \nabla_{\vec{r}}^2}{2m} \phi_R(\vec{r}) = \epsilon(R) \phi_R(\vec{r}),$$

$$\phi_R(\vec{r} \rightarrow \vec{R}_i/2) \propto \frac{1}{|\vec{r} - \vec{R}_i/2|} - \frac{1}{a}$$

2+1 (trimer) in the 3D Born-Oppenheimer picture



$$-\frac{\hbar^2 \nabla_{\vec{r}}^2}{2m} \phi_R(\vec{r}) = \epsilon(R) \phi_R(\vec{r}),$$

Light atom in the field of fixed heavy fermions

(distance $R = |\vec{R}_2 - \vec{R}_1|$)


$$\phi_R(\vec{r} \rightarrow \vec{R}_i/2) \propto \frac{1}{|\vec{r} - \vec{R}_i/2|} - \frac{1}{a}$$

Small R :

$$\epsilon_{+,m}(R) \sim -\frac{\hbar^2}{mR^2}$$

Large R :

$$\epsilon_{+,m}(R) \sim \epsilon_0, \text{ dimer energy}$$



[D. S. Petrov, arXiv:1206.5752]

2+1 (trimer) in the 3D Born-Oppenheimer picture

Schrödinger equation for heavy atom with reduced mass $M/2$ in the effective potential:


$$\left[-\frac{\hbar^2}{M} \frac{\partial^2}{\partial R^2} + U_{\text{eff}}(R) - E \right] \chi(R) = 0, \quad U_{\text{eff}}(R) = \frac{\hbar^2 l(l+1)}{MR^2} + \epsilon_{+,m}(R) + |\epsilon_0|$$

2+1 (trimer) in the 3D Born-Oppenheimer picture

Schrödinger equation for heavy atom with reduced mass $M/2$ in the effective potential:

$$\left[-\frac{\hbar^2}{M} \frac{\partial^2}{\partial R^2} + U_{\text{eff}}(R) - E \right] \chi(R) = 0, \quad U_{\text{eff}}(R) = \frac{\hbar^2 l(l+1)}{MR^2} + \epsilon_{+,m}(R) + |\epsilon_0|$$

The light-mediated effective heavy-heavy potential is “tuned” by M/m :



Outline

- ▷ Introduction and motivation
- ▷ Born-Oppenheimer theory of the 3D trimer
- ▷ Bound states of $N + 1$ fermions in 1D
- ▷ Derivation of the results
 - ▷ Exact results for $N \leq 5$
 - ▷ Mean field: Thomas-Fermi approximation
 - ▷ Mean field: Hartree-Fock
- ▷ Conclusions and perspectives

Binding in 1D


As in 3D, there is a **competition** between heavy-heavy kinetic energy and light-mediated heavy-heavy attraction.

Binding in 1D


As in 3D, there is a **competition** between **heavy-heavy kinetic energy** and **light-mediated heavy-heavy attraction**.

State of the art in 1D:

- ▶ Trimer (2+1 atoms) at $M/m \geq 1$ (red dashed curve)
[Kartavtsev, et al. JETP **108**, 365 (2009)]
- ▶ Tetramer (3+1 atoms) through Born-Oppenheimer treatment (lowest black curve)
[Mehta, PRA **89**, 052706 (2014)]



Our results




We provide the **exact**
solution of the quantum
mechanical problem up
to $N = 5$.

($N = 2, 3, 4, 5$ here)

[A. Tononi, J. Givois, and D. S. Petrov, arXiv:2205.01018]

Our results



We provide the **exact** solution of the quantum mechanical problem up to $N = 5$.

($N = 2, 3, 4, 5$ here)


We identify the critical mass ratios:

$$(M/m)_{2+1} = 1,$$

$$(M/m)_{3+1} = 1.76,$$

$$(M/m)_{4+1} = 4.2,$$

$$(M/m)_{5+1} = 12.0 \pm 0.5$$



[A. Tononi, J. Givois, and D. S. Petrov, arXiv:2205.01018]

Outline

- ▷ Introduction and motivation
- ▷ Born-Oppenheimer theory of the 3D trimer
- ▷ Bound states of $N + 1$ fermions in 1D
- ▷ Derivation of the results
 - ▷ Exact results for $N \leq 5$
 - ▷ Mean field: Thomas-Fermi approximation
 - ▷ Mean field: Hartree-Fock
- ▷ Conclusions and perspectives

Exact results: the Skorniakov-Ter Martirosian equation

Schrödinger equation for a system of N heavy plus 1 light fermions:

$$\left[-\sum_{i=1}^N \frac{\partial_{x_i}^2}{2M} - \frac{\partial_{x_{N+1}}^2}{2m} + g \sum_{i < N+1} \delta(x_i - x_{N+1}) - E \right] \psi(x_1, \dots, x_N, x_{N+1}) = 0,$$

where $E < 0$, and $g = -1/(m_r a) < 0$, $m_r = mM/(m + M)$.

Exact results: the Skorniakov-Ter Martirosian equation

Schrödinger equation for a system of N heavy plus 1 light fermions:

$$\left[-\sum_{i=1}^N \frac{\partial_{x_i}^2}{2M} - \frac{\partial_{x_{N+1}}^2}{2m} + g \sum_{i < N+1} \delta(x_i - x_{N+1}) - E \right] \psi(x_1, \dots, x_N, x_{N+1}) = 0,$$

where $E < 0$, and $g = -1/(m_r a) < 0$, $m_r = mM/(m + M)$.

Wave function of $(N - 1)$ fermions plus a dimer:

$$\psi(x_1, \dots, x_{N-1}, x_N, x_{N+1} = x_N)$$

Exact results: the Skorniakov-Ter Martirosian equation

Schrödinger equation for a system of N heavy plus 1 light fermions:

$$\left[-\sum_{i=1}^N \frac{\partial_{x_i}^2}{2M} - \frac{\partial_{x_{N+1}}^2}{2m} + g \sum_{i < N+1} \delta(x_i - x_{N+1}) - E \right] \psi(x_1, \dots, x_N, x_{N+1}) = 0,$$

where $E < 0$, and $g = -1/(m_r a) < 0$, $m_r = mM/(m + M)$.

Wave function of $(N - 1)$ fermions plus a dimer:

$$\psi(x_1, \dots, x_{N-1}, x_N, x_{N+1} = x_N)$$

Fourier transform: $F(q_1, \dots, q_{N-1}, q_N)$

Exact results: the Skorniakov-Ter Martirosian equation

Schrödinger equation for a system of N heavy plus 1 light fermions:

$$\left[-\sum_{i=1}^N \frac{\partial_{x_i}^2}{2M} - \frac{\partial_{x_{N+1}}^2}{2m} + g \sum_{i < N+1} \delta(x_i - x_{N+1}) - E \right] \psi(x_1, \dots, x_N, x_{N+1}) = 0,$$

where $E < 0$, and $g = -1/(m_r a) < 0$, $m_r = mM/(m + M)$.


Wave function of $(N - 1)$ fermions plus a dimer:

$$\psi(x_1, \dots, x_{N-1}, x_N, x_{N+1} = x_N)$$

Fourier transform: $F(q_1, \dots, q_{N-1}, q_N)$

In center of mass coordinates $q_N = -\sum_{i=1}^N q_i$ we have:

$$F(q_1, \dots, q_{N-1})$$



Exact results: the Skorniakov-Ter Martirosian equation

$F(q_1, \dots, q_{N-1})$ satisfies the STM equation

$$\left[\frac{a}{2} - \frac{1}{2\kappa(q_1, \dots, q_{N-1})} \right] F(q_1, \dots, q_{N-1}) = - \int \frac{dp}{2\pi} \frac{\sum_{j=1}^{N-1} F(q_1, \dots, q_{j-1}, p, q_{j+1}, \dots, q_{N-1})}{\kappa^2(q_1, \dots, q_{N-1}) + (p + \frac{m_r}{m} \sum_{i=1}^{N-1} q_i)^2},$$

where $\kappa(q_1, \dots, q_{N-1}) = \sqrt{-2m_r E + \frac{m_r}{M+m} (\sum_{i=1}^{N-1} q_i)^2 + \frac{m_r}{M} \sum_{i=1}^{N-1} q_i^2}$.


[Skorniakov, Ter-Martirosian, JETP **4**, 648 (1957)]

[Pricoupenko, Petrov, PRA **100**, 042707 (2019)]

Integro-differential equation that **includes naturally zero-range interactions**, and removes the dimer coordinates.

Exact results: the Skorniakov-Ter Martirosian equation

The exact solution of the STM equation gives the energies (continuous lines) of the $N + 1$ clusters:



We also find that the trimer and tetramer have $P = -1$, while pentamer and hexamer have $P = +1$

Two questions

- ▶ Large N limit?
- ▶ Are there computationally-cheap methods that work also at small N ?

Outline

- ▷ Introduction and motivation
- ▷ Born-Oppenheimer theory of the 3D trimer
- ▷ Bound states of $N + 1$ fermions in 1D
- ▷ Derivation of the results
 - ▷ Exact results for $N \leq 5$
 - ▷ Mean field: Thomas-Fermi approximation
 - ▷ Mean field: Hartree-Fock
- ▷ Conclusions and perspectives

Thomas-Fermi approach

Large N limit: mean-field theory based on the Thomas-Fermi approximation for the heavy fermions

$$\Omega = \int \left[\frac{|\phi'(x)|^2}{2m} + gn(x)|\phi(x)|^2 + \frac{\pi^2 n^3(x)}{6M} - \epsilon|\phi(x)|^2 - \mu n(x) \right] dx,$$

Thomas-Fermi approach

Large N limit: mean-field theory based on the Thomas-Fermi approximation for the heavy fermions

$$\Omega = \int \left[\frac{|\phi'(x)|^2}{2m} + gn(x)|\phi(x)|^2 + \frac{\pi^2 n^3(x)}{6M} - \epsilon|\phi(x)|^2 - \mu n(x) \right] dx,$$

minimizing Ω wrt ϕ and n : $-\phi_1''(x) - 2mgn(x)\phi_1(x) = 2m\epsilon\phi_1(x)$,

$n(x) = \sqrt{-2Mg(|\phi(x)|^2 - \mu/g)/\pi^2}$, when $|\phi(x)|^2 > \mu/g$.

Thomas-Fermi approach

Large N limit: mean-field theory based on the Thomas-Fermi approximation for the heavy fermions

$$\Omega = \int \left[\frac{|\phi'(x)|^2}{2m} + gn(x)|\phi(x)|^2 + \frac{\pi^2 n^3(x)}{6M} - \epsilon|\phi(x)|^2 - \mu n(x) \right] dx,$$

minimizing Ω wrt ϕ and n : $-\phi_1''(x) - 2mgn(x)\phi_1(x) = 2m\epsilon\phi_1(x)$,


$n(x) = \sqrt{-2Mg(|\phi(x)|^2 - \mu/g)/\pi^2}$, when $|\phi(x)|^2 > \mu/g$.

When $\mu = 0$ (threshold for binding a new heavy atom), analytical:

Threshold: $\left(\frac{M}{m}\right)_{N+1} = \frac{\pi^2}{36} N^3$

$$\phi(x) = \frac{-3\pi\epsilon}{\sqrt{-8Mg^3}} \frac{1}{\cosh^2(\sqrt{-m\epsilon/2}x)}$$

$$n(x) = \sqrt{-2Mg/\pi^2} |\phi(x)|$$




Thomas-Fermi approach

We extend the theory for $\mu \neq 0$, and calculate cluster energies.

Thomas-Fermi approach

We extend the theory for $\mu \neq 0$, and calculate cluster energies.


Thomas-Fermi approach (grey curves), **analytical**, **computationally cheap**, works at large N :



Thomas-Fermi approach

We extend the theory for $\mu \neq 0$, and calculate cluster energies.

Thomas-Fermi approach (grey curves), analytical, computationally cheap, works at large N :



What is the main source of discrepancy with the small- N exact results? TF, mean field?

Hartree-Fock approach

$$\hat{H} = \int \left(-\frac{\hat{\Psi}_x^\dagger \partial_x^2 \hat{\Psi}_x}{2M} - \frac{\hat{\phi}_x^\dagger \partial_x^2 \hat{\phi}_x}{2m} + g \hat{\Psi}_x^\dagger \hat{\phi}_x^\dagger \hat{\Psi}_x \hat{\phi}_x \right) dx$$

Energy $E_{N+1} = \langle v | \hat{H} | v \rangle$, with the variational ansatz:
 $|v\rangle = \int dx \phi_1(x) \hat{\phi}_x^\dagger \int dx_1 \dots dx_N \frac{\det[\Psi_\nu(x_\eta)]}{\sqrt{N!}} \prod_{\eta=1}^N \hat{\Psi}_{x_\eta}^\dagger |0\rangle$

Hartree-Fock approach

$$\hat{H} = \int \left(-\frac{\hat{\Psi}_x^\dagger \partial_x^2 \hat{\Psi}_x}{2M} - \frac{\hat{\phi}_x^\dagger \partial_x^2 \hat{\phi}_x}{2m} + g \hat{\Psi}_x^\dagger \hat{\phi}_x^\dagger \hat{\Psi}_x \hat{\phi}_x \right) dx$$

Energy $E_{N+1} = \langle v | \hat{H} | v \rangle$, with the variational ansatz:


$$|v\rangle = \int dx \phi_1(x) \hat{\phi}_x^\dagger \int dx_1 \dots dx_N \frac{\det[\Psi_\nu(x_\eta)]}{\sqrt{N!}} \prod_{\eta=1}^N \hat{\Psi}_{x_\eta}^\dagger |0\rangle$$

Minimizing $E_{N+1} - \epsilon_1 - \mu N$ with respect to the orbitals yields:

$$-\frac{\partial_x^2 \phi_1}{2m} + g n \phi_1 = \epsilon_1 \phi_1,$$





$$-\frac{\partial_x^2 \Psi_\nu}{2M} + g |\phi_1|^2 \Psi_\nu = E_\nu \Psi_\nu,$$

$$n = \sum_{\nu=1}^N |\Psi_\nu|^2$$







Hartree-Fock approach

→ No improvement wrt TF energies (dashed lines)



Hartree-Fock approach

→ No improvement wrt TF energies (dashed lines)




M/m

M/m

M/m

M/m



a second-order correction (dotted lines) gives good and **computationally cheap** agreement

$N - 1$ atoms momentum distribution

All methods give access to the following quantity:

$$\rho_{N+1}(q) = \int |F(q, q_2, \dots, q_{N-1})|^2 dq_2 \dots dq_{N-1},$$

that can be used to compare their effectiveness at small N .


$N - 1$ atoms momentum distribution

All methods give access to the following quantity:

$$\rho_{N+1}(q) = \int |F(q, q_2, \dots, q_{N-1})|^2 dq_2 \dots dq_{N-1},$$

that can be used to compare their effectiveness at small N .

We find that Hartree-Fock reproduces very well these momentum correlations:



Outline

- ▷ Introduction and motivation
- ▷ Born-Oppenheimer theory of the 3D trimer
- ▷ Bound states of $N + 1$ fermions in 1D
- ▷ Derivation of the results
 - ▷ Exact results for $N \leq 5$
 - ▷ Mean field: Thomas-Fermi approximation
 - ▷ Mean field: Hartree-Fock
- ▷ Conclusions and perspectives

Conclusions and perspectives

Binding of N heavy fermions by a light atom:
for larger mass ratio more atoms can be bound.

(very different from 3D, where there are no bound states for
 $M/m > 13.6$, meaning no 6+1 clusters!)

Conclusions and perspectives

Binding of N heavy fermions by a light atom:
for larger mass ratio more atoms can be bound.

(very different from 3D, where there are no bound states for
 $M/m > 13.6$, meaning no 6+1 clusters!)

- Exact results up to $N = 5$
- **TF theory**: analytical and works for large N
- **HF theory**: reproduces well energy and correlations at small and large N

Conclusions and perspectives

Binding of N heavy fermions by a light atom:
for larger mass ratio more atoms can be bound.

(very different from 3D, where there are no bound states for
 $M/m > 13.6$, meaning no 6+1 clusters!)

- Exact results up to $N = 5$
- **TF theory**: analytical and works for large N
- **HF theory**: reproduces well energy and correlations at small and large N

Possible generalization to other setups:

- higher dimensions
- more particles

Thank you for your attention!

References:

- A. Tononi, J. Givois, and D. S. Petrov, arXiv:2205.01018, accepted in PRA as a Letter
- A. Pricoupenko, and D. S. Petrov, PRA **100**, 042707 (2019)